

The Project Gutenberg EBook of Geometrical Solutions Derived from Mechanics  
Archimedes

This eBook is for the use of anyone anywhere in the United States and most other parts of the world at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at [www.gutenberg.org](http://www.gutenberg.org). If you are not located in the United States, you'll have to check the laws of the country where you are located before using this eBook.

Title: Geometrical Solutions Derived from Mechanics  
A Treatise of Archimedes

Author: Archimedes

Release Date: November 1, 2014 [EBook #7825]

Language: English

Character set encoding: ISO-8859-1

\*\*\* START OF THIS PROJECT GUTENBERG EBOOK GEOMETRICAL SOLUTIONS \*\*\*

### TRANSCRIBER'S NOTE

The camera-quality files for this public-domain ebook may be downloaded *gratis* at

[www.gutenberg.org/ebooks/7825](http://www.gutenberg.org/ebooks/7825).

This ebook was produced using scanned images generously provided by the University of California, Berkeley, Mathematics and Statistics Library, through the Internet Archive.

Minor typographical corrections and presentational changes have been made without comment.

This PDF file is optimized for screen viewing, but may be recompiled for printing. Please consult the preamble of the L<sup>A</sup>T<sub>E</sub>X source file for instructions and other particulars.

# GEOMETRICAL SOLUTIONS

DERIVED FROM  
MECHANICS

## A TREATISE OF ARCHIMEDES

RECENTLY DISCOVERED AND  
TRANSLATED FROM THE GREEK BY  
DR. J. L. HEIBERG

PROFESSOR OF CLASSICAL PHILOLOGY  
AT THE UNIVERSITY OF COPENHAGEN

WITH AN INTRODUCTION BY  
DAVID EUGENE SMITH  
PRESIDENT OF TEACHERS COLLEGE,  
COLUMBIA UNIVERSITY, NEW YORK

ENGLISH VERSION TRANSLATED FROM THE GERMAN  
BY LYDIA G. ROBINSON  
AND REPRINTED FROM "THE MONIST," APRIL, 1909

CHICAGO  
THE OPEN COURT PUBLISHING COMPANY  
LONDON AGENTS  
KEGAN PAUL, TRENCH, TRÜBNER & CO., LTD.  
1909

COPYRIGHT BY  
THE OPEN COURT PUBLISHING CO.  
1909

## INTRODUCTION.

IF there ever was a case of appropriateness in discovery, the finding of this manuscript in the summer of 1906 was one. In the first place it was appropriate that the discovery should be made in Constantinople, since it was here that the West received its first manuscripts of the other extant works, nine in number, of the great Syracusan. It was furthermore appropriate that the discovery should be made by Professor Heiberg, *facilis princeps* among all workers in the field of editing the classics of Greek mathematics, and an indefatigable searcher of the libraries of Europe for manuscripts to aid him in perfecting his labors. And finally it was most appropriate that this work should appear at a time when the affiliation of pure and applied mathematics is becoming so generally recognized all over the world. We are sometimes led to feel, in considering isolated cases, that the great contributors of the past have worked in the field of pure mathematics alone, and the saying of Plutarch that Archimedes felt that "every kind of art connected with daily needs was ignoble and vulgar"<sup>1</sup> may have strengthened this feeling. It therefore assists us in properly orientating ourselves to read another treatise from the greatest mathematician of antiquity that sets clearly before us his indebtedness to the mechanical applications of his subject.

Not the least interesting of the passages in the manuscript

---

<sup>1</sup>Marcellus, 17.

is the first line, the greeting to Eratosthenes. It is well known, on the testimony of Diodoros his countryman, that Archimedes studied in Alexandria, and the latter frequently makes mention of Konon of Samos whom he knew there, probably as a teacher, and to whom he was indebted for the suggestion of the spiral that bears his name. It is also related, this time by Proclus, that Eratosthenes was a contemporary of Archimedes, and if the testimony of so late a writer as Tzetzes, who lived in the twelfth century, may be taken as valid, the former was eleven years the junior of the great Sicilian. Until now, however, we have had nothing definite to show that the two were ever acquainted. The great Alexandrian savant,—poet, geographer, arithmetician,—affectionately called by the students Pentathlos, the champion in five sports,<sup>2</sup> selected by Ptolemy Euergetes to succeed his master, Kallimachos the poet, as head of the great Library,—this man, the most renowned of his time in Alexandria, could hardly have been a teacher of Archimedes, nor yet the fellow student of one who was so much his senior. It is more probable that they were friends in the later days when Archimedes was received as a savant rather than as a learner, and this is borne out by the statement at the close of proposition I which refers to one of his earlier works, showing that this particular treatise was a late one. This reference being to one of the two works dedicated

---

<sup>2</sup>His nickname of *Beta* is well known, possibly because his lecture room was number 2.

to Dositheos of Kolonos,<sup>3</sup> and one of these (*De lineis spiraliibus*) referring to an earlier treatise sent to Konon,<sup>4</sup> we are led to believe that this was one of the latest works of Archimedes and that Eratosthenes was a friend of his mature years, although one of long standing. The statement that the preliminary propositions were sent "some time ago" bears out this idea of a considerable duration of friendship, and the idea that more or less correspondence had resulted from this communication may be inferred by the statement that he saw, as he had previously said, that Eratosthenes was "a capable scholar and a prominent teacher of philosophy," and also that he understood "how to value a mathematical method of investigation when the opportunity offered." We have, then, new light upon the relations between these two men, the leaders among the learned of their day.

A second feature of much interest in the treatise is the intimate view that we have into the workings of the mind of the author. It must always be remembered that Archimedes was primarily a discoverer, and not primarily a compiler as were Euclid, Apollonios, and Nicomachos. Therefore to have him follow up his first communication of theorems to Eratosthenes by a statement of his mental processes in reaching his conclusions is not merely a contribution to mathematics but one to education as well. Particularly is this true in

---

<sup>3</sup>We know little of his works, none of which are extant. Geminus and Ptolemy refer to certain observations made by him in 200 B. C., twelve years after the death of Archimedes. Pliny also mentions him.

<sup>4</sup> Τῶν ποτὶ Κόνωνα ἀπυσταλέντων θεωρημάτων.

the following statement, which may well be kept in mind in the present day: "I have thought it well to analyse and lay down for you in this same book a peculiar method by means of which it will be possible for you to derive instruction as to how certain mathematical questions may be investigated by means of mechanics. And I am convinced that this is equally profitable in demonstrating a proposition itself; for much that was made evident to me through the medium of mechanics was later proved by means of geometry, because the treatment by the former method had not yet been established by way of a demonstration. For of course it is easier to establish a proof if one has in this way previously obtained a conception of the questions, than for him to seek it without such a preliminary notion. . . . Indeed I assume that some one among the investigators of to-day or in the future will discover by the method here set forth still other propositions which have not yet occurred to us." Perhaps in all the history of mathematics no such prophetic truth was ever put into words. It would almost seem as if Archimedes must have seen as in a vision the methods of Galileo, Cavalieri, Pascal, Newton, and many of the other great makers of the mathematics of the Renaissance and the present time.

The first proposition concerns the quadrature of the parabola, a subject treated at length in one of his earlier communications to Dositheos.<sup>5</sup> He gives a digest of the treatment, but with the warning that the proof is not com-

---

<sup>5</sup> Τετραγωνισμὸς παραβολῆς.



plete, as it is in his special work upon the subject. He has, in fact, summarized propositions VII–XVII of his communication to Dositheos, omitting the geometric treatment of propositions XVIII–XXIV. One thing that he does not state, here or in any of his works, is where the idea of center of gravity<sup>6</sup> started. It was certainly a common notion in his day, for he often uses it without defining it. It appears in Euclid's<sup>7</sup> time, but how much earlier we cannot as yet say.

Proposition II states no new fact. Essentially it means that if a sphere, cylinder, and cone (always circular) have the same radius,  $r$ , and the altitude of the cone is  $r$  and that of the cylinder  $2r$ , then the volumes will be as  $4 : 1 : 6$ , which is true, since they are respectively  $\frac{4}{3}\pi r^3$ ,  $\frac{1}{3}\pi r^3$ , and  $2\pi r^3$ . The interesting thing, however, is the method pursued, the derivation of geometric truths from principles of mechanics. There is, too, in every sentence, a little suggestion of Cavalieri, an anticipation by nearly two thousand years of the work of the greatest immediate precursor of Newton. And the geometric imagination that Archimedes shows in the last sentence is also noteworthy as one of the interesting features of this work: "After I had thus perceived that a sphere is four times as large as the cone. . . it occurred to me that the surface of a sphere is four times as great as its largest circle, in which I proceeded from the idea that just as a circle is

---

<sup>6</sup> Κέντρα βαρῶν, for "barycentric" is a very old term.

<sup>7</sup>At any rate in the anonymous fragment *De levi et ponderoso*, sometimes attributed to him.

equal to a triangle whose base is the periphery of the circle, and whose altitude is equal to its radius, so a sphere is equal to a cone whose base is the same as the surface of the sphere and whose altitude is equal to the radius of the sphere." As a bit of generalization this throws a good deal of light on the workings of Archimedes's mind.

In proposition III he considers the volume of a spheroid, which he had already treated more fully in one of his letters to Dositheos,<sup>8</sup> and which contains nothing new from a mathematical standpoint. Indeed it is the method rather than the conclusion that is interesting in such of the subsequent propositions as relate to mensuration. Proposition V deals with the center of gravity of a segment of a conoid, and proposition VI with the center of gravity of a hemisphere, thus carrying into solid geometry the work of Archimedes on the equilibrium of planes and on their centers of gravity.<sup>9</sup> The general method is that already known in the treatise mentioned, and this is followed through proposition X.

Proposition XI is the interesting case of a segment of a right cylinder cut off by a plane through the center of the lower base and tangent to the upper one. He shows this to equal one-sixth of the square prism that circumscribes the cylinder. This is well known to us through the formula  $v = 2r^2h/3$ , the volume of the prism being  $4r^2h$ , and requires a knowledge of the center of gravity of the cylindrical

---

<sup>8</sup> Περὶ κωνοειδῶν καὶ σφαιροειδῶν.

<sup>9</sup> Ἐπιπέδων ἰσορροπιῶν ἢ κέντρα βαρῶν ἐπιπέδων.

section in question. Archimedes is, so far as we know, the first to state this result, and he obtains it by his usual method of the skilful balancing of sections. There are several lacunae in the demonstration, but enough of it remains to show the ingenuity of the general plan. The culminating interest from the mathematical standpoint lies in proposition XIII, where Archimedes reduces the whole question to that of the quadrature of the parabola. He shows that a fourth of the circumscribed prism is to the segment of the cylinder as the semi-base of the prism is to the parabola inscribed in the semi-base; that is, that  $\frac{1}{4}p : v = \frac{1}{2}b : (\frac{2}{3} \cdot \frac{1}{2}b)$ , whence  $v = \frac{1}{6}p$ . Proposition XIV is incomplete, but it is the conclusion of the two preceding propositions.

In general, therefore, the greatest value of the work lies in the following:

1. It throws light upon the hitherto only suspected relations of Archimedes and Eratosthenes.
2. It shows the working of the mind of Archimedes in the discovery of mathematical truths, showing that he often obtained his results by intuition or even by measurement, rather than by an analytic form of reasoning, verifying these results later by strict analysis.
3. It expresses definitely the fact that Archimedes was the discoverer of those properties relating to the sphere and cylinder that have been attributed to him and that are given in his other works without a definite statement of their authorship.
4. It shows that Archimedes was the first to state the

volume of the cylinder segment mentioned, and it gives an interesting description of the mechanical method by which he arrived at his result.

DAVID EUGENE SMITH.  
TEACHERS COLLEGE, COLUMBIA UNIVERSITY.

## GEOMETRICAL SOLUTIONS DERIVED FROM MECHANICS.

ARCHIMEDES TO ERATOSTHENES, GREETING:

Some time ago I sent you some theorems I had discovered, writing down only the propositions because I wished you to find their demonstrations which had not been given. The propositions of the theorems which I sent you were the following:

1. If in a perpendicular prism with a parallelogram<sup>1</sup> for base a cylinder is inscribed which has its bases in the opposite parallelograms<sup>1</sup> and its surface touching the other planes of the prism, and if a plane is passed through the center of the circle that is the base of the cylinder and one side of the square lying in the opposite plane, then that plane will cut off from the cylinder a section which is bounded by two planes, the intersecting plane and the one in which the base of the cylinder lies, and also by as much of the surface of the cylinder as lies between these same planes; and the detached section of the cylinder is  $\frac{1}{6}$  of the whole prism.

2. If in a cube a cylinder is inscribed whose bases lie in opposite parallelograms<sup>1</sup> and whose surface touches the other four planes, and if in the same cube a second cylinder is inscribed whose bases lie in two other parallelograms<sup>1</sup> and whose surface touches the four other planes, then the body enclosed by the surface of the cylinder and comprehended

---

<sup>1</sup>This must mean a square.

within both cylinders will be equal to  $\frac{2}{3}$  of the whole cube.

These propositions differ essentially from those formerly discovered; for then we compared those bodies (conoids, spheroids and their segments) with the volume of cones and cylinders but none of them was found to be equal to a body enclosed by planes. Each of these bodies, on the other hand, which are enclosed by two planes and cylindrical surfaces is found to be equal to a body enclosed by planes. The demonstration of these propositions I am accordingly sending to you in this book.

Since I see, however, as I have previously said, that you are a capable scholar and a prominent teacher of philosophy, and also that you understand how to value a mathematical method of investigation when the opportunity is offered, I have thought it well to analyze and lay down for you in this same book a peculiar method by means of which it will be possible for you to derive instruction as to how certain mathematical questions may be investigated by means of mechanics. And I am convinced that this is equally profitable in demonstrating a proposition itself; for much that was made evident to me through the medium of mechanics was later proved by means of geometry because the treatment by the former method had not yet been established by way of a demonstration. For of course it is easier to establish a proof if one has in this way previously obtained a conception of the questions, than for him to seek it without such a preliminary notion. Thus in the familiar propositions the demonstrations of which Eudoxos was the first to discover,

namely that a cone and a pyramid are one third the size of that cylinder and prism respectively that have the same base and altitude, no little credit is due to Democritos who was the first to make that statement about these bodies without any demonstration. But we are in a position to have found the present proposition in the same way as the earlier one; and I have decided to write down and make known the method partly because we have already talked about it heretofore and so no one would think that we were spreading abroad idle talk, and partly in the conviction that by this means we are obtaining no slight advantage for mathematics, for indeed I assume that some one among the investigators of to-day or in the future will discover by the method here set forth still other propositions which have not yet occurred to us.

In the first place we will now explain what was also first made clear to us through mechanics, namely that a segment of a parabola is  $\frac{4}{3}$  of the triangle possessing the same base and equal altitude; following which we will explain in order the particular propositions discovered by the above mentioned method; and in the last part of the book we will present the geometrical demonstrations of the propositions.<sup>2</sup>

1. If one magnitude is taken away from another magnitude and the same point is the center of gravity both of the

---

<sup>2</sup>In his "Commentar," Professor Zeuthen calls attention to the fact that it was already known from Heron's recently discovered *Metrica* that these propositions were contained in this treatise, and Professor Heiberg made the same comment in *Hermes*.—Tr.

whole and of the part removed, then the same point is the center of gravity of the remaining portion.

2. If one magnitude is taken away from another magnitude and the center of gravity of the whole and of the part removed is not the same point, the center of gravity of the remaining portion may be found by prolonging the straight line which connects the centers of gravity of the whole and of the part removed, and setting off upon it another straight line which bears the same ratio to the straight line between the aforesaid centers of gravity, as the weight of the magnitude which has been taken away bears to the weight of the one remaining [*De plan. aequil.* I, 8].

3. If the centers of gravity of any number of magnitudes lie upon the same straight line, then will the center of gravity of all the magnitudes combined lie also upon the same straight line [Cf. *ibid.* I, 5].

4. The center of gravity of a straight line is the center of that line [Cf. *ibid.* I, 4].

5. The center of gravity of a triangle is the point in which the straight lines drawn from the angles of a triangle to the centers of the opposite sides intersect [*Ibid.* I, 14].

6. The center of gravity of a parallelogram is the point where its diagonals meet [*Ibid.* I, 10].

7. The center of gravity [of a circle] is the center [of that circle].

8. The center of gravity of a cylinder [is the center of its axis].

9. The center of gravity of a prism is the center of its



axis.

10. The center of gravity of a cone so divides its axis that the section at the vertex is three times as great as the remainder.

11. Moreover together with the exercise here laid down I will make use of the following proposition:

If any number of magnitudes stand in the same ratio to the same number of other magnitudes which correspond pair by pair, and if either all or some of the former magnitudes stand in any ratio whatever to other magnitudes, and the latter in the same ratio to the corresponding ones, then the sum of the magnitudes of the first series will bear the same ratio to the sum of those taken from the third series as the sum of those of the second series bears to the sum of those taken from the fourth series [*De Conoid.* I].

## I.

Let  $\alpha\beta\gamma$  [Fig. 1] be the segment of a parabola bounded by the straight line  $\alpha\gamma$  and the parabola  $\alpha\beta\gamma$ . Let  $\alpha\gamma$  be bisected at  $\delta$ ,  $\delta\beta\epsilon$  being parallel to the diameter, and draw  $\alpha\beta$ , and  $\beta\gamma$ . Then the segment  $\alpha\beta\gamma$  will be  $\frac{4}{3}$  as great as the triangle  $\alpha\beta\gamma$ .

From the points  $\alpha$  and  $\gamma$  draw  $\alpha\zeta \parallel \delta\beta\epsilon$ , and the tangent  $\gamma\zeta$ ; produce  $[\gamma\beta$  to  $\kappa$ , and make  $\kappa\theta = \gamma\kappa$ ]. Think of  $\gamma\theta$  as a scale-beam with the center at  $\kappa$  and let  $\mu\xi$  be any straight line whatever  $\parallel \epsilon\delta$ . Now since  $\gamma\beta\alpha$  is a parabola,  $\gamma\zeta$  a tangent and  $\gamma\delta$  an ordinate, then  $\epsilon\beta = \beta\delta$ ; for this

indeed has been proved in the Elements [i. e., of conic sections, cf. *Quadr. parab.* 2]. For this reason and because  $\zeta\alpha$  and  $\mu\xi \parallel \epsilon\delta$ ,  $\mu\nu = \nu\xi$ , and  $\zeta\kappa = \kappa\alpha$ . And because  $\gamma\alpha : \alpha\xi = \mu\xi : \xi o$  (for this is shown in a corollary, [cf. *Quadr. parab.* 5]),  $\gamma\alpha : \alpha\xi = \gamma\kappa : \kappa\nu$ ; and  $\gamma\kappa = \kappa\theta$ , therefore  $\theta\kappa : \kappa\nu = \mu\xi : \xi o$ . And because  $\nu$  is the center of gravity of the straight line  $\mu\xi$ , since  $\mu\nu = \nu\xi$ , then if we make  $\tau\eta = \xi o$  and  $\theta$  as its center of gravity so that  $\tau\theta = \theta\eta$ , the straight line  $\tau\theta\eta$  will be in equilibrium with  $\mu\xi$  in its present position because  $\theta\nu$  is divided in inverse proportion to the weights  $\tau\eta$  and  $\mu\xi$ , and  $\theta\kappa : \kappa\nu = \mu\xi : \eta\tau$ ; therefore  $\kappa$  is the center of gravity of the combined weight of the two. In the




Fig. 1.

same way all straight lines drawn in the triangle  $\zeta\alpha\gamma \parallel \epsilon\delta$

are in their present positions in equilibrium with their parts cut off by the parabola, when these are transferred to  $\theta$ , so that  $\kappa$  is the center of gravity of the combined weight of the two. And because the triangle  $\gamma\zeta\alpha$  consists of the straight lines in the triangle  $\gamma\zeta\alpha$  and the segment  $\alpha\beta\gamma$  consists of those straight lines within the segment of the parabola corresponding to the straight line  $\xi o$ , therefore the triangle  $\zeta\alpha\gamma$  in its present position will be in equilibrium at the point  $\kappa$  with the parabola-segment when this is transferred to  $\theta$  as its center of gravity, so that  $\kappa$  is the center of gravity of the combined weights of the two. Now let  $\gamma\kappa$  be so divided at  $\chi$  that  $\gamma\kappa = 3\kappa\chi$ ; then  $\chi$  will be the center of gravity of the triangle  $\alpha\zeta\gamma$ , for this has been shown in the Statics [cf. *De plan. aequil.* I, 15, p. 186, 3 with Eutokios, S. 320, 5ff.]. Now the triangle  $\zeta\alpha\gamma$  in its present position is in equilibrium at the point  $\kappa$  with the segment  $\beta\alpha\gamma$  when this is transferred to  $\theta$  as its center of gravity, and the center of gravity of the triangle  $\zeta\alpha\gamma$  is  $\chi$ ; hence triangle  $\alpha\zeta\gamma$  : segm.  $\alpha\beta\gamma$  when transferred to  $\theta$  as its center of gravity =  $\theta\kappa$  :  $\kappa\chi$ . But  $\theta\kappa = 3\kappa\chi$ ; hence also triangle  $\alpha\zeta\gamma = 3$  segm.  $\alpha\beta\gamma$ . But it is also true that triangle  $\zeta\alpha\gamma = 4\Delta\alpha\beta\gamma$  because  $\zeta\kappa = \kappa\alpha$  and  $\alpha\delta = \delta\gamma$ ; hence segm.  $\alpha\beta\gamma = \frac{4}{3}$  the triangle  $\alpha\beta\gamma$ . This is of course clear.

It is true that this is not proved by what we have said here; but it indicates that the result is correct. And so, as we have just seen that it has not been proved but rather conjectured that the result is correct we have devised a geometrical demonstration which we made known some time

ago and will again bring forward farther on.

## II.

That a sphere is four times as large as a cone whose base is equal to the largest circle of the sphere and whose altitude is equal to the radius of the sphere, and that a cylinder whose base is equal to the largest circle of the sphere and whose altitude is equal to the diameter of the circle is one and a half times as large as the sphere, may be seen by the present method in the following way:




Fig. 2.

Let  $\alpha\beta\gamma\delta$  [Fig. 2] be the largest circle of a sphere and

$\alpha\gamma$  and  $\beta\delta$  its diameters perpendicular to each other; let there be in the sphere a circle on the diameter  $\beta\delta$  perpendicular to the circle  $\alpha\beta\gamma\delta$ , and on this perpendicular circle let there be a cone erected with its vertex at  $\alpha$ ; producing the convex surface of the cone, let it be cut through  $\gamma$  by a plane parallel to its base; the result will be the circle perpendicular to  $\alpha\gamma$  whose diameter will be  $\epsilon\zeta$ . On this circle erect a cylinder whose axis =  $\alpha\gamma$  and whose vertical boundaries are  $\epsilon\lambda$  and  $\zeta\eta$ . Produce  $\gamma\alpha$  making  $\alpha\theta = \gamma\alpha$  and think of  $\gamma\theta$  as a scale-beam with its center at  $\alpha$ . Then let  $\mu\nu$  be any straight line whatever drawn  $\parallel \beta\delta$  intersecting the circle  $\alpha\beta\gamma\delta$  in  $\xi$  and  $o$ , the diameter  $\alpha\gamma$  in  $\sigma$ , the straight line  $\alpha\epsilon$  in  $\pi$  and  $\alpha\zeta$  in  $\rho$ , and on the straight line  $\mu\nu$  construct a plane perpendicular to  $\alpha\gamma$ ; it will intersect the cylinder in a circle on the diameter  $\mu\nu$ ; the sphere  $\alpha\beta\gamma\delta$ , in a circle on the diameter  $\xi o$ ; the cone  $\alpha\epsilon\zeta$  in a circle on the diameter  $\pi\rho$ . Now because  $\gamma\alpha \times \alpha\sigma = \mu\sigma \times \sigma\pi$  (for  $\alpha\gamma = \sigma\mu$ ,  $\alpha\sigma = \pi\sigma$ ), and  $\gamma\alpha \times \alpha\sigma = \alpha\xi^2 = \xi\sigma^2 + \alpha\pi^2$  then  $\mu\sigma \times \sigma\pi = \xi\sigma^2 + \sigma\pi^2$ . Moreover, because  $\gamma\alpha : \alpha\sigma = \mu\sigma : \sigma\pi$  and  $\gamma\alpha = \alpha\theta$ , therefore  $\theta\alpha : \alpha\sigma = \mu\sigma : \sigma\pi = \mu\sigma^2 : \mu\sigma \times \sigma\pi$ . But it has been proved that  $\xi\sigma^2 + \sigma\pi^2 = \mu\sigma \times \sigma\pi$ ; hence  $\alpha\theta : \alpha\sigma = \mu\sigma^2 : \xi\sigma^2 + \sigma\pi^2$ . But it is true that  $\mu\sigma^2 : \xi\sigma^2 + \sigma\pi^2 = \mu\nu^2 : \xi\alpha^2 + \pi\rho^2 =$  the circle in the cylinder whose diameter is  $\mu\nu$  : the circle in the cone whose diameter is  $\pi\rho$  + the circle in the sphere whose diameter is  $\xi o$ ; hence  $\theta\alpha : \alpha\sigma =$  the circle in the cylinder : the circle in the sphere + the circle in the cone. Therefore the circle in the cylinder in its present position will be in equilibrium at the point  $\alpha$  with the two circles whose di-

ameters are  $\xi o$  and  $\pi\rho$ , if they are so transferred to  $\theta$  that  $\theta$  is the center of gravity of both. In the same way it can be shown that when another straight line is drawn in the parallelogram  $\xi\lambda \parallel \epsilon\zeta$ , and upon it a plane is erected perpendicular to  $\alpha\gamma$ , the circle produced in the cylinder in its present position will be in equilibrium at the point  $\alpha$  with the two circles produced in the sphere and the cone when they are transferred and so arranged on the scale-beam at the point  $\theta$  that  $\theta$  is the center of gravity of both. Therefore if cylinder, sphere and cone are filled up with such circles then the cylinder in its present position will be in equilibrium at the point  $\alpha$  with the sphere and the cone together, if they are transferred and so arranged on the scale-beam at the point  $\theta$  that  $\theta$  is the center of gravity of both. Now since the bodies we have mentioned are in equilibrium, the cylinder with  $\kappa$  as its center of gravity, the sphere and the cone transferred as we have said so that they have  $\theta$  as center of gravity, then  $\theta\alpha : \alpha\kappa = \text{cylinder} : \text{sphere} + \text{cone}$ . But  $\theta\alpha = 2\alpha\kappa$ , and hence also the cylinder =  $2 \times (\text{sphere} + \text{cone})$ . But it is also true that the cylinder = 3 cones [Euclid, *Elem.* XII, 10], hence 3 cones = 2 cones + 2 spheres. If 2 cones be subtracted from both sides, then the cone whose axes form the triangle  $\alpha\epsilon\zeta = 2$  spheres. But the cone whose axes form the triangle  $\alpha\epsilon\zeta = 8$  cones whose axes form the triangle  $\alpha\beta\delta$  because  $\epsilon\zeta = 2\beta\delta$ , hence the aforesaid 8 cones = 2 spheres. Consequently the sphere whose greatest circle is  $\alpha\beta\gamma\delta$  is four times as large as the cone with its vertex at  $\alpha$ , and whose base is the circle on the diameter  $\beta\delta$  perpendicular to  $\alpha\gamma$ .

Draw the straight lines  $\phi\beta\chi$  and  $\psi\delta\omega \parallel \alpha\gamma$  through  $\beta$  and  $\delta$  in the parallelogram  $\lambda\zeta$  and imagine a cylinder whose bases are the circles on the diameters  $\phi\psi$  and  $\chi\omega$  and whose axis is  $\alpha\gamma$ . Now since the cylinder whose axes form the parallelogram  $\phi\omega$  is twice as large as the cylinder whose axes form the parallelogram  $\phi\delta$  and the latter is three times as large as the cone the triangle of whose axes is  $\alpha\beta\delta$ , as is shown in the Elements [Euclid, *Elem.* XII, 10], the cylinder whose axes form the parallelogram  $\phi\omega$  is six times as large as the cone whose axes form the triangle  $\alpha\beta\delta$ . But it was shown that the sphere whose largest circle is  $\alpha\beta\gamma\delta$  is four times as large as the same cone, consequently the cylinder is one and one half times as large as the sphere, Q. E. D.

After I had thus perceived that a sphere is four times as large as the cone whose base is the largest circle of the sphere and whose altitude is equal to its radius, it occurred to me that the surface of a sphere is four times as great as its largest circle, in which I proceeded from the idea that just as a circle is equal to a triangle whose base is the periphery of the circle and whose altitude is equal to its radius, so a sphere is equal to a cone whose base is the same as the surface of the sphere and whose altitude is equal to the radius of the sphere.

### III.

By this method it may also be seen that a cylinder whose base is equal to the largest circle of a spheroid and whose altitude is equal to the axis of the spheroid, is one and one half

times as large as the spheroid, and when this is recognized it becomes clear that if a spheroid is cut through its center by a plane perpendicular to its axis, one-half of the spheroid is twice as great as the cone whose base is that of the segment and its axis the same.

For let a spheroid be cut by a plane through its axis and let there be in its surface an ellipse  $\alpha\beta\gamma\delta$  [Fig. 3] whose diameters are  $\alpha\gamma$  and  $\beta\delta$  and whose center is  $\kappa$  and let there be a circle in the spheroid on the diameter  $\beta\delta$  perpendicular to  $\alpha\gamma$ ; then imagine a cone whose base is the same circle but whose vertex is at  $\alpha$ , and producing its surface, let the cone be cut by a plane through  $\gamma$  parallel to the base; the intersection will be a circle perpendicular to  $\alpha\gamma$  with  $\epsilon\zeta$  as its diameter. Now imagine a cylinder whose base is the same circle with the diameter  $\epsilon\zeta$  and whose axis is  $\alpha\gamma$ ; let  $\gamma\alpha$  be produced so that  $\alpha\theta = \gamma\alpha$ ; think of  $\theta\gamma$  as a scale-beam with its center at  $\alpha$  and in the parallelogram  $\lambda\theta$  draw a straight line  $\mu\nu \parallel \epsilon\zeta$ , and on  $\mu\nu$  construct a plane perpendicular to  $\alpha\gamma$ ; this will intersect the cylinder in a circle whose diameter is  $\mu\nu$ , the spheroid in a circle whose diameter is  $\xi\sigma$  and the cone in a circle whose diameter is  $\pi\rho$ . Because  $\gamma\alpha : \alpha\sigma = \epsilon\alpha : \alpha\pi = \mu\sigma : \sigma\pi$ , and  $\gamma\alpha = \alpha\theta$ , therefore  $\theta\alpha : \alpha\sigma = \mu\sigma : \sigma\pi$ . But  $\mu\sigma : \sigma\pi = \mu\sigma^2 : \mu\sigma \times \sigma\pi$  and  $\mu\sigma \times \sigma\pi = \pi\sigma^2 + \sigma\xi^2$ , for  $\alpha\sigma \times \sigma\gamma : \sigma\xi^2 = \alpha\kappa \times \kappa\gamma : \kappa\beta^2 = \alpha\kappa^2 : \kappa\beta^2$  (for both ratios are equal to the ratio between the diameter and the parameter [Apollonius, *Con.* I, 21]) =  $\alpha\sigma^2 : \sigma\pi^2$  therefore  $\alpha\sigma^2 : \alpha\sigma \times \sigma\gamma = \pi\sigma^2 : \sigma\xi^2 = \sigma\pi^2 : \sigma\pi \times \pi\mu$ , consequently  $\mu\pi \times \pi\sigma = \sigma\xi^2$ . If  $\pi\sigma^2$  is added to both sides then



$\mu\sigma \times \sigma\pi = \pi\sigma^2 + \sigma\xi^2$ . Therefore  $\theta\alpha : \alpha\sigma = \mu\sigma^2 : \pi\sigma^2 + \sigma\xi^2$ .  
 But  $\mu\sigma^2 : \sigma\xi^2 + \sigma\pi^2 =$  the circle in the cylinder whose diameter is  $\mu\nu$  : the circle with the diameter  $\xi o$  + the circle with the diameter  $\pi\rho$ ; hence the circle whose diameter is  $\mu\nu$  will in its present position be in equilibrium at the point  $\alpha$  with the two circles whose diameters are  $\xi o$  and  $\pi\rho$  when they are transferred and so arranged on the scale-beam at the point  $\alpha$  that  $\theta$  is the center of gravity of both; and  $\theta$  is the center of gravity of the two circles combined whose diameters are  $\xi o$  and  $\pi\rho$  when their position is changed, hence  $\theta\alpha : \alpha\sigma =$  the




Fig. 3.

circle with the diameter  $\mu\nu$  : the two circles whose diameters are  $\xi o$  and  $\pi\rho$ . In the same way it can be shown that if another straight line is drawn in the parallelogram  $\lambda\zeta \parallel \epsilon\zeta$  and on this line last drawn a plane is constructed perpendicular to  $\alpha\gamma$ , then likewise the circle produced in the cylinder will in its present position be in equilibrium at the point  $\alpha$  with the two circles combined which have been produced in the spheroid and in the cone respectively when they are so transferred to the point  $\theta$  on the scale-beam that  $\theta$  is the center of gravity of both. Then if cylinder, spheroid and cone are filled with such circles, the cylinder in its present position will be in equilibrium at the point  $\alpha$  with the spheroid + the cone if they are transferred and so arranged on the scale-beam at the point  $\alpha$  that  $\theta$  is the center of gravity of both. Now  $\kappa$  is the center of gravity of the cylinder, but  $\theta$ , as has been said, is the center of gravity of the spheroid and cone together. Therefore  $\theta\alpha : \alpha\kappa = \text{cylinder} : \text{spheroid} + \text{cone}$ . But  $\alpha\theta = 2\alpha\kappa$ , hence also the cylinder =  $2 \times (\text{spheroid} + \text{cone}) = 2 \times \text{spheroid} + 2 \times \text{cone}$ . But the cylinder =  $3 \times \text{cone}$ , hence  $3 \times \text{cone} = 2 \times \text{cone} + 2 \times \text{spheroid}$ . Subtract  $2 \times \text{cone}$  from both sides; then a cone whose axes form the triangle  $\alpha\epsilon\zeta = 2 \times \text{spheroid}$ . But the same cone = 8 cones whose axes form the  $\Delta\alpha\beta\delta$ ; hence 8 such cones =  $2 \times \text{spheroid}$ ,  $4 \times \text{cone} = \text{spheroid}$ ; whence it follows that a spheroid is four times as great as a cone whose vertex is at  $\alpha$ , and whose base is the circle on the diameter  $\beta\delta$  perpendicular to  $\lambda\epsilon$ , and one-half the spheroid is twice as great as the same cone.

In the parallelogram  $\lambda\zeta$  draw the straight lines  $\phi\chi$  and

$\psi\omega \parallel \alpha\gamma$  through the points  $\beta$  and  $\delta$  and imagine a cylinder whose bases are the circles on the diameters  $\phi\psi$  and  $\chi\omega$ , and whose axis is  $\alpha\gamma$ . Now since the cylinder whose axes form the parallelogram  $\phi\omega$  is twice as great as the cylinder whose axes form the parallelogram  $\phi\delta$  because their bases are equal but the axis of the first is twice as great as the axis of the second, and since the cylinder whose axes form the parallelogram  $\phi\delta$  is three times as great as the cone whose vertex is at  $\alpha$  and whose base is the circle on the diameter  $\beta\delta$  perpendicular to  $\alpha\gamma$ , then the cylinder whose axes form the parallelogram  $\phi\omega$  is six times as great as the aforesaid cone. But it has been shown that the spheroid is four times as great as the same cone, hence the cylinder is one and one half times as great as the spheroid. Q. E. D.

#### IV.

That a segment of a right conoid cut by a plane perpendicular to its axis is one and one half times as great as the cone having the same base and axis as the segment, can be proved by the same method in the following way:

Let a right conoid be cut through its axis by a plane intersecting the surface in a parabola  $\alpha\beta\gamma$  [Fig. 4]; let it be also cut by another plane perpendicular to the axis, and let their common line of intersection be  $\beta\gamma$ . Let the axis of the segment be  $\delta\alpha$  and let it be produced to  $\theta$  so that  $\theta\alpha = \alpha\delta$ . Now imagine  $\delta\theta$  to be a scale-beam with its center at  $\alpha$ ; let the base of the segment be the circle on the diameter  $\beta\gamma$

perpendicular to  $\alpha\delta$ ; imagine a cone whose base is the circle on the diameter  $\beta\gamma$ , and whose vertex is at  $\alpha$ . Imagine also a cylinder whose base is the circle on the diameter  $\beta\gamma$  and its axis  $\alpha\delta$ , and in the parallelogram let a straight line  $\mu\nu$  be drawn  $\parallel \beta\gamma$  and on  $\mu\nu$  construct a plane perpendicular to  $\alpha\delta$ ; it will intersect the cylinder in a circle whose diameter is  $\mu\nu$ , and the segment of the right conoid in a circle whose diameter is  $\xi o$ . Now since  $\beta\alpha\gamma$  is a parabola,  $\alpha\delta$  its diameter and  $\xi\sigma$  and  $\beta\delta$  its ordinates, then [*Quadr. parab.* 3]  $\delta\alpha : \alpha\sigma = \beta\delta^2 : \xi\sigma^2$ . But  $\delta\alpha = \alpha\theta$ , therefore  $\theta\alpha : \alpha\sigma = \mu\sigma^2 : \sigma\xi^2$ . But  $\mu\sigma^2 : \sigma\xi^2 =$  the circle in the cylinder whose diameter is  $\mu\nu$  : the circle in the segment of the right conoid whose diameter is  $\xi o$ , hence  $\theta\alpha : \alpha\sigma =$  the circle with the diameter  $\mu\nu$  : the circle with the diameter  $\xi o$ ; therefore the circle in the cylinder whose diameter is  $\mu\nu$  is in its present position, in equilibrium at the point  $\alpha$  with the circle whose diameter is  $\xi o$  if this be transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. And the center of gravity of the circle whose diameter is  $\mu\nu$  is at  $\sigma$ , that of the circle whose diameter is  $\xi o$  when its position is changed, is  $\theta$ , and we have the inverse proportion,  $\theta\alpha : \alpha\sigma =$  the circle with the diameter  $\mu\nu$  : the circle with the diameter  $\xi o$ . In the same way it can be shown that if another straight line be drawn in the parallelogram  $\epsilon\gamma \parallel \beta\gamma$  the circle formed in the cylinder, will in its present position be in equilibrium at the point  $\alpha$  with that formed in the segment of the right conoid if the latter is so transferred to  $\theta$  on the scale-beam that  $\theta$  is its center of gravity. Therefore if the cylinder and the




Fig. 4.

segment of the right conoid are filled up then the cylinder in its present position will be in equilibrium at the point  $\alpha$  with the segment of the right conoid if the latter is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. And since these magnitudes are in equilibrium at  $\alpha$ , and  $\kappa$  is the center of gravity of the cylinder, if  $\alpha\delta$  is bisected at  $\kappa$  and  $\theta$  is the center of gravity of the segment transferred to that point, then we have the inverse proportion  $\theta\alpha : \alpha\kappa = \text{cylinder} : \text{segment}$ . But  $\theta\alpha = 2\alpha\kappa$  and also the cylinder =  $2 \times$  segment. But the same cylinder is 3 times as great as the cone whose base is the circle on the diameter  $\beta\gamma$  and whose vertex is at  $\alpha$ ; therefore it is clear that the segment

is one and one half times as great as the same cone.

## V.

That the center of gravity of a segment of a right conoid which is cut off by a plane perpendicular to the axis, lies on the straight line which is the axis of the segment divided in such a way that the portion at the vertex is twice as great as the remainder, may be perceived by our method in the following way:

Let a segment of a right conoid cut off by a plane perpendicular to the axis be cut by another plane through the axis, and let the intersection in its surface be the parabola  $\alpha\beta\gamma$  [Fig. 5] and let the common line of intersection of the plane which cut off the segment and of the intersecting plane be  $\beta\gamma$ ; let the axis of the segment and the diameter of the parabola  $\alpha\beta\gamma$  be  $\alpha\delta$ ; produce  $\delta\alpha$  so that  $\alpha\theta = \alpha\delta$  and imagine  $\delta\theta$  to be a scale-beam with its center at  $\alpha$ ; then inscribe a cone in the segment with the lateral boundaries  $\beta\alpha$  and  $\alpha\gamma$  and in the parabola draw a straight line  $\xi o \parallel \beta\gamma$  and let it cut the parabola in  $\xi$  and  $o$  and the lateral boundaries of the cone in  $\pi$  and  $\rho$ . Now because  $\xi\sigma$  and  $\beta\delta$  are drawn perpendicular to the diameter of the parabola,  $\delta\alpha : \alpha\sigma = \beta\delta^2 : \xi\sigma^2$  [*Quadr. parabol.* 3]. But  $\delta\alpha : \alpha\sigma = \beta\delta : \pi\sigma = \beta\delta^2 : \beta\delta \times \pi\sigma$ , therefore also  $\beta\delta^2 : \xi\sigma^2 = \beta\delta^2 : \beta\delta \times \pi\sigma$ . Consequently  $\xi\sigma^2 = \beta\delta \times \pi\sigma$  and  $\beta\delta : \xi\sigma = \xi\sigma : \pi\sigma$ , therefore  $\beta\delta : \pi\sigma = \xi\sigma^2 : \sigma\pi^2$ . But  $\beta\delta : \pi\sigma = \delta\alpha : \alpha\sigma = \theta\alpha : \alpha\sigma$ , therefore also  $\theta\alpha : \alpha\sigma = \xi\sigma^2 : \sigma\pi^2$ . On  $\xi o$  construct a plane perpen-




Fig. 5.

dicular to  $\alpha\delta$ ; this will intersect the segment of the right conoid in a circle whose diameter is  $\xi o$  and the cone in a circle whose diameter is  $\pi\rho$ . Now because  $\theta\alpha : \alpha\sigma = \xi\sigma^2 : \sigma\pi^2$  and  $\xi\sigma^2 : \sigma\pi^2 =$  the circle with the diameter  $\xi o$  : the circle with the diameter  $\pi\rho$ , therefore  $\theta\alpha : \alpha\sigma =$  the circle whose diameter is  $\xi o$  : the circle whose diameter is  $\pi\rho$ . Therefore the circle whose diameter is  $\xi o$  will in its present position be in equilibrium at the point  $\alpha$  with the circle whose diameter is  $\pi\rho$  when this is so transferred to  $\theta$  on the scale-beam that  $\theta$  is its center of gravity. Now since  $\sigma$  is the center of gravity

of the circle whose diameter is  $\xi o$  in its present position, and  $\theta$  is the center of gravity of the circle whose diameter is  $\pi\rho$  if its position is changed as we have said, and inversely  $\theta\alpha : \alpha\sigma =$  the circle with the diameter  $\xi o$  : the circle with the diameter  $\pi\rho$ , then the circles are in equilibrium at the point  $\alpha$ . In the same way it can be shown that if another straight line is drawn in the parabola  $\parallel \beta\gamma$  and on this line last drawn a plane is constructed perpendicular to  $\alpha\delta$ , the circle formed in the segment of the right conoid will in its present position be in equilibrium at the point  $\alpha$  with the circle formed in the cone, if the latter is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. Therefore if the segment and the cone are filled up with circles, all circles in the segment will be in their present positions in equilibrium at the point  $\alpha$  with all circles of the cone if the latter are transferred and so arranged on the scale-beam at the point  $\theta$  that  $\theta$  is their center of gravity. Therefore also the segment of the right conoid in its present position will be in equilibrium at the point  $\alpha$  with the cone if it is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. Now because the center of gravity of both magnitudes taken together is  $\alpha$ , but that of the cone alone when its position is changed is  $\theta$ , then the center of gravity of the remaining magnitude lies on  $\alpha\theta$  extended towards  $\alpha$  if  $\alpha\kappa$  is cut off in such a way that  $\alpha\theta : \alpha\kappa =$  segment : cone. But the segment is one and one half the size of the cone, consequently  $\alpha\theta = \frac{3}{2}\alpha\kappa$  and  $\kappa$ , the center of gravity of the right conoid, so divides  $\alpha\delta$  that the



portion at the vertex of the segment is twice as large as the remainder.

## VI.

[The center of gravity of a hemisphere is so divided on its axis] that the portion near the surface of the hemisphere is in the ratio of 5 : 3 to the remaining portion.

Let a sphere be cut by a plane through its center intersecting the surface in the circle  $\alpha\beta\gamma\delta$  [Fig. 6],  $\alpha\gamma$  and  $\beta\delta$  being two diameters of the circle perpendicular to each other. Let a plane be constructed on  $\beta\delta$  perpendicular to  $\alpha\gamma$ . Then imagine a cone whose base is the circle with the diameter  $\beta\delta$ , whose vertex is at  $\alpha$  and its lateral boundaries are  $\beta\alpha$  and  $\alpha\delta$ ; let  $\gamma\alpha$  be produced so that  $\alpha\theta = \gamma\alpha$ , imagine the straight line  $\theta\gamma$  to be a scale-beam with its center at  $\alpha$  and in the semicircle  $\beta\alpha\delta$  draw a straight line  $\xi o \parallel \beta\delta$ ; let it cut the circumference of the semicircle in  $\xi$  and  $o$ , the lateral boundaries of the cone in  $\pi$  and  $\rho$ , and  $\alpha\gamma$  in  $\epsilon$ . On  $\xi o$  construct a plane perpendicular to  $\alpha\epsilon$ ; it will intersect the hemisphere in a circle with the diameter  $\xi o$ , and the cone in a circle with the diameter  $\pi\rho$ . Now because  $\alpha\gamma : \alpha\epsilon = \xi\alpha^2 : \alpha\epsilon^2$  and  $\xi\alpha^2 = \alpha\epsilon^2 + \epsilon\xi^2$  and  $\alpha\epsilon = \epsilon\pi$ , therefore  $\alpha\gamma : \alpha\epsilon = \xi\epsilon^2 + \epsilon\pi^2 : \epsilon\pi^2$ . But  $\xi\epsilon^2 + \epsilon\pi^2 : \epsilon\pi^2 =$  the circle with the diameter  $\xi o +$  the circle with the diameter  $\pi\rho :$  the circle with the diameter  $\pi\rho$ , and  $\gamma\alpha = \alpha\theta$ , hence  $\theta\alpha : \alpha\epsilon =$  the circle with the diameter  $\xi o +$  the circle with the diameter  $\pi\rho :$  circle with the diameter  $\pi\rho$ . Therefore the




Fig. 6.

two circles whose diameters are  $\xi o$  and  $\pi\rho$  in their present position are in equilibrium at the point  $\alpha$  with the circle whose diameter is  $\pi\rho$  if it is transferred and so arranged at  $\theta$  that  $\theta$  is its center of gravity. Now since the center of gravity of the two circles whose diameters are  $\xi o$  and  $\pi\rho$  in

their present position [is the point  $\epsilon$ , but of the circle whose diameter is  $\pi\rho$  when its position is changed is the point  $\theta$ , then  $\theta\alpha : \alpha\epsilon =$  the circles whose diameters are]  $\xi o$ ,  $\pi\rho$  : the circle whose diameter is  $\pi\rho$ . In the same way if another straight line in the] hemisphere  $\beta\alpha\delta$  [is drawn  $\parallel \beta\delta$  and a plane is constructed] perpendicular to [ $\alpha\gamma$  the] two [circles produced in the cone and in the hemisphere are in their position] in equilibrium at  $\alpha$  [with the circle which is produced in the cone] if it is transferred and arranged on the scale at  $\theta$ . [Now if] the hemisphere and the cone [are filled up with circles then all circles in the] hemisphere and those [in the cone] will in their present position be in equilibrium [with all circles] in the cone, if these are transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is their center of gravity; [therefore the hemisphere and cone also] are in their position [in equilibrium at the point  $\alpha$ ] with the cone if it is transferred and so arranged [on the scale-beam at  $\theta$ ] that  $\theta$  is its center of gravity.

## VII.

By [this method] it may also be perceived that [any segment whatever] of a sphere bears the same ratio to a cone having the same [base] and axis [that the radius of the sphere + the axis of the opposite segment : the axis of the opposite segment] .....  
 and [Fig. 7] on  $\mu\nu$  construct a plane perpendicular to  $\alpha\gamma$ ; it will intersect the cylinder in a circle whose diameter is  $\mu\nu$ ,

the segment of the sphere in a circle whose diameter is  $\xi o$




Fig. 7.

and the cone whose base is the circle on the diameter  $\epsilon\zeta$  and whose vertex is at  $\alpha$  in a circle whose diameter is  $\pi\rho$ . In the same way as before it may be shown that a circle whose diameter is  $\mu\nu$  is in its present position in equilibrium at  $\alpha$  with the two circles [whose diameters are  $\xi o$  and  $\pi\rho$  if they are so arranged on the scale-beam that  $\theta$  is their center of gravity. [And the same can be proved of all corresponding circles.] Now since cylinder, cone, and spherical segment are filled up with such circles, the cylinder in its present position [will be in equilibrium at  $\alpha$ ] with the cone + the

spherical segment if they are transferred and attached to the scale-beam at  $\theta$ . Divide  $\alpha\eta$  at  $\phi$  and  $\chi$  so that  $\alpha\chi = \chi\eta$  and  $\eta\phi = \frac{1}{3}\alpha\phi$ ; then  $\chi$  will be the center of gravity of the cylinder because it is the center of the axis  $\alpha\eta$ . Now because the above mentioned bodies are in equilibrium at  $\alpha$ , cylinder : cone with the diameter of its base  $\epsilon\zeta$  + the spherical segment  $\beta\alpha\delta = \theta\alpha : \alpha\chi$ . And because  $\eta\alpha = 3\eta\phi$  then  $[\gamma\eta \times \eta\phi] = \frac{1}{3}\alpha\eta \times \eta\gamma$ . Therefore also  $\gamma\eta \times \eta\phi = \frac{1}{3}\beta\eta^2$ . . . . .

VIIa.

In the same way it may be perceived that any segment of an ellipsoid cut off by a perpendicular plane, bears the same ratio to a cone having the same base and the same axis, as half of the axis of the ellipsoid + the axis of the opposite segment bears to the axis of the opposite segment. . . . .

VIII.

.....  
 produce  $\alpha\gamma$  [Fig. 8] making  $\alpha\theta = \alpha\gamma$  and  $\gamma\xi =$  the radius of the sphere; imagine  $\gamma\theta$  to be a scale-beam with a center at  $\alpha$ , and in the plane cutting off the segment inscribe a circle with its center at  $\eta$  and its radius =  $\alpha\eta$ ; on this circle construct a cone with its vertex at  $\alpha$  and its lateral boundaries  $\alpha\epsilon$  and  $\alpha\zeta$ . Then draw a straight line  $\kappa\lambda \parallel \epsilon\zeta$ ; let it cut the circumference of the segment at  $\kappa$  and  $\lambda$ , the lateral boundaries of the cone  $\alpha\epsilon\zeta$  at  $\rho$  and  $o$  and  $\alpha\gamma$  at  $\pi$ . Now because




Fig. 8.

$\alpha\gamma : \alpha\pi = \alpha\kappa^2 : \alpha\pi^2$  and  $\kappa\alpha^2 = \alpha\pi^2 + \pi\kappa^2$  and  $\alpha\pi^2 = \pi o^2$   
 (since also  $\alpha\eta^2 = \epsilon\eta^2$ ), then  $\gamma\alpha : \alpha\pi = \kappa\pi^2 + \pi o^2 : \pi o^2$ . But  
 $\kappa\pi^2 + \pi o^2 : \pi o^2 =$  the circle with the diameter  $\kappa\lambda +$  the circle with the diameter  $o\rho$  : the circle with the diameter  $o\rho$  and  
 $\gamma\alpha = \alpha\theta$ ; therefore  $\theta\alpha : \alpha\pi =$  the circle with the diameter  $\kappa\lambda +$  the circle with the diameter  $o\rho$  : the circle with the diameter  $o\rho$ . Now since the circle with the diameter  $\kappa\lambda +$  the circle with the diameter  $o\rho$  : the circle with the diam-

eter  $o\rho = \alpha\theta : \pi\alpha$ , let the circle with the diameter  $o\rho$  be transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity; then  $\theta\alpha : \alpha\pi =$  the circle with the diameter  $\kappa\lambda +$  the circle with the diameter  $o\rho$  in their present positions : the circle with the diameter  $o\rho$  if it is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. Therefore the circles in the segment  $\beta\alpha\delta$  and in the cone  $\alpha\epsilon\zeta$  are in equilibrium at  $\alpha$  with that in the cone  $\alpha\epsilon\zeta$ . And in the same way all circles in the segment  $\beta\alpha\delta$  and in the cone  $\alpha\epsilon\zeta$  in their present positions are in equilibrium at the point  $\alpha$  with all circles in the cone  $\alpha\epsilon\zeta$  if they are transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is their center of gravity; then also the spherical segment  $\alpha\beta\delta$  and the cone  $\alpha\epsilon\zeta$  in their present positions are in equilibrium at the point  $\alpha$  with the cone  $\epsilon\alpha\zeta$  if it is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity. Let the cylinder  $\mu\nu$  equal the cone whose base is the circle with the diameter  $\epsilon\zeta$  and whose vertex is at  $\alpha$  and let  $\alpha\eta$  be so divided at  $\phi$  that  $\alpha\eta = 4\phi\eta$ ; then  $\phi$  is the center of gravity of the cone  $\epsilon\alpha\zeta$  as has been previously proved. Moreover let the cylinder  $\mu\nu$  be so cut by a perpendicularly intersecting plane that the cylinder  $\mu$  is in equilibrium with the cone  $\epsilon\alpha\zeta$ . Now since the segment  $\alpha\beta\delta +$  the cone  $\epsilon\alpha\zeta$  in their present positions are in equilibrium at  $\alpha$  with the cone  $\epsilon\alpha\zeta$  if it is transferred and so arranged on the scale-beam at  $\theta$  that  $\theta$  is its center of gravity, and cylinder  $\mu\nu =$  cone  $\epsilon\alpha\zeta$  and the two cylinders  $\mu + \nu$  are moved to  $\theta$  and  $\mu\nu$  is in equilibrium with both bodies, then will also the cylinder  $\nu$  be in equilibrium

with the segment of the sphere at the point  $\alpha$ . And since the spherical segment  $\beta\alpha\delta$  : the cone whose base is the circle with the diameter  $\beta\delta$ , and whose vertex is at  $\alpha = \xi\eta : \eta\gamma$  (for this has previously been proved [*De sph. et cyl.* II, 2 Coroll.]) and cone  $\beta\alpha\delta$  : cone  $\epsilon\alpha\zeta =$  the circle with the diameter  $\beta\delta$  : the circle with the diameter  $\epsilon\zeta = \beta\eta^2 : \eta\epsilon^2$ , and  $\beta\eta^2 = \gamma\eta \times \eta\alpha$ ,  $\eta\epsilon^2 = \eta\alpha^2$ , and  $\gamma\eta \times \eta\alpha : \eta\alpha^2 = \gamma\eta : \eta\alpha$ , therefore cone  $\beta\alpha\delta$  : cone  $\epsilon\alpha\zeta = \gamma\eta : \eta\alpha$ . But we have shown that cone  $\beta\alpha\delta$  : segment  $\beta\alpha\delta = \gamma\eta : \eta\xi$ , hence  $\delta\acute{\iota}\ \acute{\iota}\sigma\omicron\upsilon$  segment  $\beta\alpha\delta$  : cone  $\epsilon\alpha\zeta = \xi\eta : \eta\alpha$ . And because  $\alpha\chi : \chi\eta = \eta\alpha + 4\eta\gamma : \alpha\eta + 2\eta\gamma$  so inversely  $\eta\chi : \chi\alpha = 2\gamma\eta + \eta\alpha : 4\gamma\eta + \eta\alpha$  and by addition  $\eta\alpha : \alpha\chi = 6\gamma\eta + 2\eta\alpha : \eta\alpha + 4\eta\gamma$ . But  $\eta\xi = \frac{1}{4}(6\eta\gamma + 2\eta\alpha)$  and  $\gamma\phi = \frac{1}{4}(4\eta\gamma + \eta\alpha)$ ; for that is evident. Hence  $\eta\alpha : \alpha\chi = \xi\eta : \gamma\phi$ , consequently also  $\xi\eta : \eta\alpha = \gamma\phi : \chi\alpha$ . But it was also demonstrated that  $\xi\eta : \eta\alpha =$  the segment whose vertex is at  $\alpha$  and whose base is the circle with the diameter  $\beta\delta$  : the cone whose vertex is at  $\alpha$  and whose base is the circle with the diameter  $\epsilon\zeta$ ; hence segment  $\beta\alpha\delta$  : cone  $\epsilon\alpha\zeta = \gamma\phi : \chi\alpha$ . And since the cylinder  $\mu$  is in equilibrium with the cone  $\epsilon\alpha\zeta$  at  $\alpha$ , and  $\theta$  is the center of gravity of the cylinder while  $\phi$  is that of the cone  $\epsilon\alpha\zeta$ , then cone  $\epsilon\alpha\zeta$  : cylinder  $\mu = \theta\alpha : \alpha\phi = \gamma\alpha : \alpha\phi$ . But cylinder  $\mu\nu =$  cone  $\epsilon\alpha\zeta$ ; hence by subtraction, cylinder  $\mu$  : cylinder  $\nu = \alpha\phi : \gamma\phi$ . And cylinder  $\mu\nu =$  cone  $\epsilon\alpha\zeta$ ; hence cone  $\epsilon\alpha\zeta$  : cylinder  $\nu = \gamma\alpha : \gamma\phi = \theta\alpha : \gamma\phi$ . But it was also demonstrated that segment  $\beta\alpha\delta$  : cone  $\epsilon\alpha\zeta = \gamma\phi : \chi\alpha$ ; hence  $\delta\acute{\iota}\ \acute{\iota}\sigma\omicron\upsilon$  segment  $\beta\alpha\delta$  : cylinder  $\nu = \zeta\alpha : \alpha\chi$ . And it was demonstrated that segment  $\beta\alpha\delta$  is in equilibrium at  $\alpha$



with the cylinder  $\nu$  and  $\theta$  is the center of gravity of the cylinder  $\nu$ , consequently the point  $\chi$  is also the center of gravity of the segment  $\beta\alpha\delta$ .

## IX.

In a similar way it can also be perceived that the center of gravity of any segment of an ellipsoid lies on the straight line which is the axis of the segment so divided that the portion at the vertex of the segment bears the same ratio to the remaining portion as the axis of the segment + 4 times the axis of the opposite segment bears to the axis of the segment + twice the axis of the opposite segment.

## X.

It can also be seen by this method that [a segment of a hyperboloid] bears the same ratio to a cone having the same base and axis as the segment, that the axis of the segment + 3 times the addition to the axis bears to the axis of the segment of the hyperboloid + twice its addition [*De Conoid.* 25]; and that the center of gravity of the hyperboloid so divides the axis that the part at the vertex bears the same ratio to the rest that three times the axis + eight times the addition to the axis bears to the axis of the hyperboloid + 4 times the addition to the axis, and many other points which I will leave aside since the method has been made clear by the examples already given and only the



demonstrations of the above given theorems remain to be stated.

## XI.

When in a perpendicular prism with square bases a cylinder is inscribed whose bases lie in opposite squares and whose curved surface touches the four other parallelograms, and when a plane is passed through the center of the circle which is the base of the cylinder and one side of the opposite square, then the body which is cut off by this plane [from the cylinder] will be  $\frac{1}{6}$  of the entire prism. This can be perceived through the present method and when it is so warranted we will pass over to the geometrical proof of it.

Imagine a perpendicular prism with square bases and a cylinder inscribed in the prism in the way we have described. Let the prism be cut through the axis by a plane perpendicular to the plane which cuts off the section of the cylinder; this will intersect the prism containing the cylinder in the parallelogram  $\alpha\beta$  [Fig. 9] and the common intersecting line of the plane which cuts off the section of the cylinder and the plane lying through the axis perpendicular to the one cutting off the section of the cylinder will be  $\beta\gamma$ ; let the axis of the cylinder and the prism be  $\gamma\delta$  which is bisected at right angles by  $\epsilon\zeta$  and on  $\epsilon\zeta$  let a plane be constructed perpendicular to  $\gamma\delta$ . This will intersect the prism in a square and the cylinder in a circle.

Now let the intersection of the prism be the square  $\mu\nu$



[Fig. 10], that of the cylinder, the circle  $\xi o \pi \rho$  and let the circle touch the sides of the square at the points  $\xi, o, \pi$  and  $\rho$ ; let the common line of intersection of the plane cutting off the cylinder-section and that passing through  $\epsilon\zeta$  perpendicular to the axis of the cylinder, be  $\kappa\lambda$ ; this line is bisected by  $\pi\theta\xi$ . In the semicircle  $o\pi\rho$  draw a straight line  $\sigma\tau$  perpendicular to  $\pi\chi$ , on  $\sigma\tau$  construct a plane perpendicular to  $\xi\pi$  and produce it to both sides of the plane enclosing the circle  $\xi o \pi \rho$ ; this will intersect the half-cylinder whose base is the semicircle  $o\pi\rho$  and whose altitude is the axis of the prism, in a parallelogram one side of which =  $\sigma\tau$  and the other = the vertical boundary of the cylinder, and it will intersect the cylinder-section likewise in a parallelogram of which one side is  $\sigma\tau$  and the other  $\mu\nu$  [Fig. 9]; and accordingly  $\mu\nu$  will be drawn in the parallelogram  $\delta\epsilon \parallel \beta\omega$  and will cut off  $\epsilon\iota = \pi\chi$ . Now because  $\epsilon\gamma$  is a parallelogram and  $\nu\iota \parallel \theta\gamma$ , and  $\epsilon\theta$  and  $\beta\gamma$  cut the parallels, therefore  $\epsilon\theta : \theta\iota = \omega\gamma : \gamma\nu = \beta\omega : \nu\nu$ .

But  $\beta\omega : \nu\nu =$  parallelogram in the half-cylinder : parallelogram in the cylinder-section, therefore both parallelograms have the same side  $\sigma\tau$ ; and  $\epsilon\theta = \theta\pi$ ,  $\iota\theta = \chi\theta$ ; and since  $\pi\theta = \theta\xi$  therefore  $\theta\xi : \theta\chi =$  parallelogram in half-cylinder : parallelogram in the cylinder-section. Imagine the parallelogram in the cylinder-section transferred and so brought to  $\xi$  that  $\xi$  is its center of gravity, and further imagine  $\pi\xi$  to be a scale-beam with its center at  $\theta$ ; then the parallelogram in the half-cylinder in its present position is in equilibrium at the point  $\theta$  with the parallelogram in the cylinder-section when it is transferred and so arranged on the scale-beam at  $\xi$  that  $\xi$  is its center of gravity. And since  $\chi$  is the center of gravity in the parallelogram in the half-cylinder, and  $\xi$  that of the parallelogram in the cylinder-section when its position is changed, and  $\xi\theta : \theta\chi =$  the parallelogram whose center of gravity is  $\chi$  : the parallelogram whose center of gravity is  $\xi$ , then the parallelogram whose center of gravity is  $\chi$  will be in equilibrium at  $\theta$  with the parallelogram whose center of gravity is  $\xi$ . In this way it can be proved that if another straight line is drawn in the semicircle  $\sigma\pi\rho$  perpendicular to  $\pi\theta$  and on this straight line a plane is constructed perpendicular to  $\pi\theta$  and is produced towards both sides of the plane in which the circle  $\xi\sigma\pi\rho$  lies, then the parallelogram formed in the half-cylinder in its present position will be in equilibrium at the point  $\theta$  with the parallelogram formed in the cylinder-section if this is transferred and so arranged on the scale-beam at  $\xi$  that  $\xi$  is its center of gravity; therefore also all parallelograms in the half-cylinder in their present

positions will be in equilibrium at the point  $\theta$  with all parallelograms of the cylinder-section if they are transferred and attached to the scale-beam at the point  $\xi$ ; consequently also the half-cylinder in its present position will be in equilibrium at the point  $\theta$  with the cylinder-section if it is transferred and so arranged on the scale-beam at  $\xi$  that  $\xi$  is its center of gravity.

## XII.

Let the parallelogram  $\mu\nu$  be perpendicular to the axis [of the circle]  $\xi o$  [ $\pi\rho$ ] [Fig. 11]. Draw  $\theta\mu$  and  $\theta\eta$  and erect upon them two planes perpendicular to the plane in which the semicircle  $o\pi\rho$  lies and extend these planes on both sides. The result is a prism whose base is a triangle similar to  $\theta\mu\eta$  and whose altitude is equal to the axis of the cylinder, and this prism is  $\frac{1}{4}$  of the entire prism which contains the cylinder. In the semicircle  $o\pi\rho$  and in the square  $\mu\nu$  draw two straight lines  $\kappa\lambda$  and  $\tau\nu$  at equal distances from  $\pi\xi$ ; these will cut the circumference of the semicircle  $o\pi\rho$  at the points  $\kappa$  and  $\tau$ , the diameter  $o\rho$  at  $\sigma$  and  $\zeta$  and the straight lines  $\theta\eta$  and  $\theta\mu$  at  $\phi$  and  $\chi$ . Upon  $\kappa\lambda$  and  $\tau\nu$  construct two planes perpendicular to  $o\rho$  and extend them towards both sides of the plane in which lies the circle  $\xi o\pi\rho$ ; they will intersect the half-cylinder whose base is the semicircle  $o\pi\rho$  and whose altitude is that of the cylinder, in a parallelogram one side of which =  $\kappa\sigma$  and the other = the axis of the cylinder; and they will intersect the prism  $\theta\eta\mu$  likewise in a parallelogram




Fig. 11.

one side of which is equal to  $\lambda\chi$  and the other equal to the axis, and in the same way the half-cylinder in a parallelogram one side of which =  $\tau\zeta$  and the other = the axis of the cylinder, and the prism in a parallelogram one side of which =  $\nu\phi$  and the other = the axis of the cylinder. ....

XIII.

Let the square  $\alpha\beta\gamma\delta$  [Fig. 12] be the base of a perpendicular prism with square bases and let a cylinder be inscribed in the prism whose base is the circle  $\epsilon\zeta\eta\theta$  which touches the sides of the parallelogram  $\alpha\beta\gamma\delta$  at  $\epsilon$ ,  $\zeta$ ,  $\eta$ , and  $\theta$ . Pass a

plane through its center and the side in the square opposite the square  $\alpha\beta\gamma\delta$  corresponding to the side  $\gamma\delta$ ; this will cut off from the whole prism a second prism which is  $\frac{1}{4}$  the size of the whole prism and which will be bounded by three parallelograms and two opposite triangles. In the semicircle  $\epsilon\zeta\eta$  describe a parabola whose origin is  $\eta\epsilon$  and whose axis is  $\zeta\kappa$ , and in the parallelogram  $\delta\eta$  draw  $\mu\nu \parallel \kappa\zeta$ ; this will cut the circumference of the semicircle at  $\xi$ , the parabola at  $\lambda$ , and  $\mu\nu \times \nu\lambda = \nu\zeta^2$  (for this is evident [Apollonios, *Con.* I, 11]). Therefore  $\mu\nu : \nu\lambda = \kappa\eta^2 : \lambda\sigma^2$ . Upon  $\mu\nu$  construct a plane parallel to  $\epsilon\eta$ ; this will intersect the prism cut off from the whole prism in a right-angled triangle one side of which is  $\mu\nu$  and the other a straight line in the plane upon  $\gamma\delta$  perpendicular to  $\gamma\delta$  at  $\nu$  and equal to the axis of the cylinder, but whose hypotenuse is in the intersecting plane. It will intersect the portion which is cut off from the cylinder by the plane passed through  $\epsilon\eta$  and the side of the square opposite the side  $\gamma\delta$  in a right-angled triangle one side of which is  $\mu\xi$  and the other a straight line drawn in the surface of the cylinder perpendicular to the plane  $\kappa\nu$ , and the hypotenuse .....

and all the triangles in the prism : all the triangles in the cylinder-section = all the straight lines in the parallelogram  $\delta\eta$  : all the straight lines between the parabola and the straight line  $\epsilon\eta$ . And the prism consists of the triangles in the prism, the cylinder-section of those in the cylinder-section, the parallelogram  $\delta\eta$  of the straight lines in the parallelogram  $\delta\eta \parallel \kappa\zeta$  and the segment of the parabola of the straight




Fig. 12.

lines cut off by the parabola and the straight line  $\epsilon\eta$ ; hence prism : cylinder-section = parallelogram  $\eta\delta$  : segment  $\epsilon\zeta\eta$  that is bounded by the parabola and the straight line  $\epsilon\eta$ . But the parallelogram  $\delta\eta = \frac{3}{2}$  the segment bounded by the parabola and the straight line  $\epsilon\eta$  as indeed has been shown in the previously published work, hence also the prism is equal to one and one half times the cylinder-section. Therefore when the cylinder-section = 2, the prism = 3 and the whole prism containing the cylinder equals 12, because it is four times the size of the other prism; hence the cylinder-section is equal to  $\frac{1}{6}$  of the prism, Q. E. D.



XIV.

[Inscribe a cylinder in] a perpendicular prism with square bases [and let it be cut by a plane passed through the center of the base of the cylinder and one side of the opposite square.] Then this plane will cut off a prism from the whole prism and a portion of the cylinder from the cylinder. It may be proved that the portion cut off from the cylinder by the plane is one-sixth of the whole prism. But first we will prove that it is possible to inscribe a solid figure in the cylinder-section and to circumscribe another composed of prisms of equal altitude and with similar triangles as bases, so that the circumscribed figure exceeds the inscribed less than any given magnitude. ....

But it has been shown that the prism cut off by the inclined plane  $< \frac{3}{2}$  the body inscribed in the cylinder-section. Now the prism cut off by the inclined plane : the body inscribed in the cylinder-section = parallelogram  $\delta\eta$  : the parallelograms which are inscribed in the segment bounded by the parabola and the straight line  $\epsilon\eta$ . Hence the parallelogram  $\delta\eta < \frac{3}{2}$  the parallelograms in the segment bounded by the parabola and the straight line  $\epsilon\eta$ . But this is impossible because we have shown elsewhere that the parallelogram  $\delta\eta$  is one and one half times the segment bounded by the parabola and the straight line  $\epsilon\eta$ , consequently is ..... not greater .....

And all prisms in the prism cut off by the inclined plane : all prisms in the figure described around the cylinder-

section = all parallelograms in the parallelogram  $\delta\eta$  : all  
 parallelograms in the figure which is described around the  
 segment bounded by the parabola and the straight line  $\epsilon\eta$ ,  
 i. e., the prism cut off by the inclined plane : the figure  
 described around the cylinder-section = parallelogram  $\delta\eta$  :  
 the figure bounded by the parabola and the straight line  $\epsilon\eta$ .  
 But the prism cut off by the inclined plane is greater than  
 one and one half times the solid figure circumscribed around  
 the cylinder-section .....

End of the Project Gutenberg EBook of Geometrical Solutions Derived from  
Mechanics, by Archimedes

\*\*\* END OF THIS PROJECT GUTENBERG EBOOK GEOMETRICAL SOLUTIONS \*\*\*

\*\*\*\*\* This file should be named 7825-pdf.pdf or 7825-pdf.zip \*\*\*\*\*  
This and all associated files of various formats will be found in:  
<http://www.gutenberg.org/7/8/2/7825/>

Produced by Gordon Keener

Updated editions will replace the previous one--the old editions  
will be renamed.

Creating the works from public domain print editions means that no  
one owns a United States copyright in these works, so the Foundation  
(and you!) can copy and distribute it in the United States without  
permission and without paying copyright royalties. Special rules,  
set forth in the General Terms of Use part of this license, apply to  
copying and distributing Project Gutenberg-tm electronic works to  
protect the PROJECT GUTENBERG-tm concept and trademark. Project  
Gutenberg is a registered trademark, and may not be used if you  
charge for the eBooks, unless you receive specific permission. If you  
do not charge anything for copies of this eBook, complying with the  
rules is very easy. You may use this eBook for nearly any purpose  
such as creation of derivative works, reports, performances and  
research. They may be modified and printed and given away--you may do  
practically ANYTHING with public domain eBooks. Redistribution is  
subject to the trademark license, especially commercial  
redistribution.

\*\*\* START: FULL LICENSE \*\*\*

THE FULL PROJECT GUTENBERG LICENSE  
PLEASE READ THIS BEFORE YOU DISTRIBUTE OR USE THIS WORK

To protect the Project Gutenberg-tm mission of promoting the free  
distribution of electronic works, by using or distributing this work

(or any other work associated in any way with the phrase "Project Gutenberg"), you agree to comply with all the terms of the Full Project Gutenberg-tm License available with this file or online at [www.gutenberg.org/license](http://www.gutenberg.org/license).

## Section 1. General Terms of Use and Redistributing Project Gutenberg-tm electronic works

1.A. By reading or using any part of this Project Gutenberg-tm electronic work, you indicate that you have read, understand, agree to and accept all the terms of this license and intellectual property (trademark/copyright) agreement. If you do not agree to abide by all the terms of this agreement, you must cease using and return or destroy all copies of Project Gutenberg-tm electronic works in your possession. If you paid a fee for obtaining a copy of or access to a Project Gutenberg-tm electronic work and you do not agree to be bound by the terms of this agreement, you may obtain a refund from the person or entity to whom you paid the fee as set forth in paragraph 1.E.8.

1.B. "Project Gutenberg" is a registered trademark. It may only be used on or associated in any way with an electronic work by people who agree to be bound by the terms of this agreement. There are a few things that you can do with most Project Gutenberg-tm electronic works even without complying with the full terms of this agreement. See paragraph 1.C below. There are a lot of things you can do with Project Gutenberg-tm electronic works if you follow the terms of this agreement and help preserve free future access to Project Gutenberg-tm electronic works. See paragraph 1.E below.

1.C. The Project Gutenberg Literary Archive Foundation ("the Foundation" or PGLAF), owns a compilation copyright in the collection of Project Gutenberg-tm electronic works. Nearly all the individual works in the collection are in the public domain in the United States. If an individual work is in the public domain in the United States and you are located in the United States, we do not claim a right to prevent you from copying, distributing, performing, displaying or creating derivative works based on the work as long as all references to Project Gutenberg are removed. Of course, we hope that you will support the Project Gutenberg-tm mission of promoting free access to electronic works by

freely sharing Project Gutenberg-tm works in compliance with the terms of this agreement for keeping the Project Gutenberg-tm name associated with the work. You can easily comply with the terms of this agreement by keeping this work in the same format with its attached full Project Gutenberg-tm License when you share it without charge with others.

1.D. The copyright laws of the place where you are located also govern what you can do with this work. Copyright laws in most countries are in a constant state of change. If you are outside the United States, check the laws of your country in addition to the terms of this agreement before downloading, copying, displaying, performing, distributing or creating derivative works based on this work or any other Project Gutenberg-tm work. The Foundation makes no representations concerning the copyright status of any work in any country outside the United States.

1.E. Unless you have removed all references to Project Gutenberg:

1.E.1. The following sentence, with active links to, or other immediate access to, the full Project Gutenberg-tm License must appear prominently whenever any copy of a Project Gutenberg-tm work (any work on which the phrase "Project Gutenberg" appears, or with which the phrase "Project Gutenberg" is associated) is accessed, displayed, performed, viewed, copied or distributed:

This eBook is for the use of anyone anywhere at no cost and with almost no restrictions whatsoever. You may copy it, give it away or re-use it under the terms of the Project Gutenberg License included with this eBook or online at [www.gutenberg.org](http://www.gutenberg.org)

1.E.2. If an individual Project Gutenberg-tm electronic work is derived from the public domain (does not contain a notice indicating that it is posted with permission of the copyright holder), the work can be copied and distributed to anyone in the United States without paying any fees or charges. If you are redistributing or providing access to a work with the phrase "Project Gutenberg" associated with or appearing on the work, you must comply either with the requirements of paragraphs 1.E.1 through 1.E.7 or obtain permission for the use of the work and the Project Gutenberg-tm trademark as set forth in paragraphs 1.E.8 or 1.E.9.

1.E.3. If an individual Project Gutenberg-tm electronic work is posted with the permission of the copyright holder, your use and distribution must comply with both paragraphs 1.E.1 through 1.E.7 and any additional terms imposed by the copyright holder. Additional terms will be linked to the Project Gutenberg-tm License for all works posted with the permission of the copyright holder found at the beginning of this work.

1.E.4. Do not unlink or detach or remove the full Project Gutenberg-tm License terms from this work, or any files containing a part of this work or any other work associated with Project Gutenberg-tm.

1.E.5. Do not copy, display, perform, distribute or redistribute this electronic work, or any part of this electronic work, without prominently displaying the sentence set forth in paragraph 1.E.1 with active links or immediate access to the full terms of the Project Gutenberg-tm License.

1.E.6. You may convert to and distribute this work in any binary, compressed, marked up, nonproprietary or proprietary form, including any word processing or hypertext form. However, if you provide access to or distribute copies of a Project Gutenberg-tm work in a format other than "Plain Vanilla ASCII" or other format used in the official version posted on the official Project Gutenberg-tm web site ([www.gutenberg.org](http://www.gutenberg.org)), you must, at no additional cost, fee or expense to the user, provide a copy, a means of exporting a copy, or a means of obtaining a copy upon request, of the work in its original "Plain Vanilla ASCII" or other form. Any alternate format must include the full Project Gutenberg-tm License as specified in paragraph 1.E.1.

1.E.7. Do not charge a fee for access to, viewing, displaying, performing, copying or distributing any Project Gutenberg-tm works unless you comply with paragraph 1.E.8 or 1.E.9.

1.E.8. You may charge a reasonable fee for copies of or providing access to or distributing Project Gutenberg-tm electronic works provided that

- You pay a royalty fee of 20% of the gross profits you derive from the use of Project Gutenberg-tm works calculated using the method

you already use to calculate your applicable taxes. The fee is owed to the owner of the Project Gutenberg-tm trademark, but he has agreed to donate royalties under this paragraph to the Project Gutenberg Literary Archive Foundation. Royalty payments must be paid within 60 days following each date on which you prepare (or are legally required to prepare) your periodic tax returns. Royalty payments should be clearly marked as such and sent to the Project Gutenberg Literary Archive Foundation at the address specified in Section 4, "Information about donations to the Project Gutenberg Literary Archive Foundation."

- You provide a full refund of any money paid by a user who notifies you in writing (or by e-mail) within 30 days of receipt that s/he does not agree to the terms of the full Project Gutenberg-tm License. You must require such a user to return or destroy all copies of the works possessed in a physical medium and discontinue all use of and all access to other copies of Project Gutenberg-tm works.
- You provide, in accordance with paragraph 1.F.3, a full refund of any money paid for a work or a replacement copy, if a defect in the electronic work is discovered and reported to you within 90 days of receipt of the work.
- You comply with all other terms of this agreement for free distribution of Project Gutenberg-tm works.

1.E.9. If you wish to charge a fee or distribute a Project Gutenberg-tm electronic work or group of works on different terms than are set forth in this agreement, you must obtain permission in writing from both the Project Gutenberg Literary Archive Foundation and Michael Hart, the owner of the Project Gutenberg-tm trademark. Contact the Foundation as set forth in Section 3 below.

#### 1.F.

1.F.1. Project Gutenberg volunteers and employees expend considerable effort to identify, do copyright research on, transcribe and proofread public domain works in creating the Project Gutenberg-tm collection. Despite these efforts, Project Gutenberg-tm electronic

works, and the medium on which they may be stored, may contain "Defects," such as, but not limited to, incomplete, inaccurate or corrupt data, transcription errors, a copyright or other intellectual property infringement, a defective or damaged disk or other medium, a computer virus, or computer codes that damage or cannot be read by your equipment.

1.F.2. LIMITED WARRANTY, DISCLAIMER OF DAMAGES - Except for the "Right of Replacement or Refund" described in paragraph 1.F.3, the Project Gutenberg Literary Archive Foundation, the owner of the Project Gutenberg-tm trademark, and any other party distributing a Project Gutenberg-tm electronic work under this agreement, disclaim all liability to you for damages, costs and expenses, including legal fees. YOU AGREE THAT YOU HAVE NO REMEDIES FOR NEGLIGENCE, STRICT LIABILITY, BREACH OF WARRANTY OR BREACH OF CONTRACT EXCEPT THOSE PROVIDED IN PARAGRAPH 1.F.3. YOU AGREE THAT THE FOUNDATION, THE TRADEMARK OWNER, AND ANY DISTRIBUTOR UNDER THIS AGREEMENT WILL NOT BE LIABLE TO YOU FOR ACTUAL, DIRECT, INDIRECT, CONSEQUENTIAL, PUNITIVE OR INCIDENTAL DAMAGES EVEN IF YOU GIVE NOTICE OF THE POSSIBILITY OF SUCH DAMAGE.

1.F.3. LIMITED RIGHT OF REPLACEMENT OR REFUND - If you discover a defect in this electronic work within 90 days of receiving it, you can receive a refund of the money (if any) you paid for it by sending a written explanation to the person you received the work from. If you received the work on a physical medium, you must return the medium with your written explanation. The person or entity that provided you with the defective work may elect to provide a replacement copy in lieu of a refund. If you received the work electronically, the person or entity providing it to you may choose to give you a second opportunity to receive the work electronically in lieu of a refund. If the second copy is also defective, you may demand a refund in writing without further opportunities to fix the problem.

1.F.4. Except for the limited right of replacement or refund set forth in paragraph 1.F.3, this work is provided to you 'AS-IS', WITH NO OTHER WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR ANY PURPOSE.

1.F.5. Some states do not allow disclaimers of certain implied



warranties or the exclusion or limitation of certain types of damages. If any disclaimer or limitation set forth in this agreement violates the law of the state applicable to this agreement, the agreement shall be interpreted to make the maximum disclaimer or limitation permitted by the applicable state law. The invalidity or unenforceability of any provision of this agreement shall not void the remaining provisions.

1.F.6. INDEMNITY - You agree to indemnify and hold the Foundation, the trademark owner, any agent or employee of the Foundation, anyone providing copies of Project Gutenberg-tm electronic works in accordance with this agreement, and any volunteers associated with the production, promotion and distribution of Project Gutenberg-tm electronic works, harmless from all liability, costs and expenses, including legal fees, that arise directly or indirectly from any of the following which you do or cause to occur: (a) distribution of this or any Project Gutenberg-tm work, (b) alteration, modification, or additions or deletions to any Project Gutenberg-tm work, and (c) any Defect you cause.

## Section 2. Information about the Mission of Project Gutenberg-tm

Project Gutenberg-tm is synonymous with the free distribution of electronic works in formats readable by the widest variety of computers including obsolete, old, middle-aged and new computers. It exists because of the efforts of hundreds of volunteers and donations from people in all walks of life.

Volunteers and financial support to provide volunteers with the assistance they need are critical to reaching Project Gutenberg-tm's goals and ensuring that the Project Gutenberg-tm collection will remain freely available for generations to come. In 2001, the Project Gutenberg Literary Archive Foundation was created to provide a secure and permanent future for Project Gutenberg-tm and future generations. To learn more about the Project Gutenberg Literary Archive Foundation and how your efforts and donations can help, see Sections 3 and 4 and the Foundation information page at [www.gutenberg.org](http://www.gutenberg.org)

## Section 3. Information about the Project Gutenberg Literary Archive Foundation

The Project Gutenberg Literary Archive Foundation is a non profit 501(c)(3) educational corporation organized under the laws of the state of Mississippi and granted tax exempt status by the Internal Revenue Service. The Foundation's EIN or federal tax identification number is 64-6221541. Contributions to the Project Gutenberg Literary Archive Foundation are tax deductible to the full extent permitted by U.S. federal laws and your state's laws.

The Foundation's principal office is located at 4557 Melan Dr. S. Fairbanks, AK, 99712., but its volunteers and employees are scattered throughout numerous locations. Its business office is located at 809 North 1500 West, Salt Lake City, UT 84116, (801) 596-1887. Email contact links and up to date contact information can be found at the Foundation's web site and official page at [www.gutenberg.org/contact](http://www.gutenberg.org/contact)

For additional contact information:

Dr. Gregory B. Newby  
Chief Executive and Director  
[gnewby@pglaf.org](mailto:gnewby@pglaf.org)

#### Section 4. Information about Donations to the Project Gutenberg Literary Archive Foundation

Project Gutenberg-tm depends upon and cannot survive without wide spread public support and donations to carry out its mission of increasing the number of public domain and licensed works that can be freely distributed in machine readable form accessible by the widest array of equipment including outdated equipment. Many small donations (\$1 to \$5,000) are particularly important to maintaining tax exempt status with the IRS.

The Foundation is committed to complying with the laws regulating charities and charitable donations in all 50 states of the United States. Compliance requirements are not uniform and it takes a considerable effort, much paperwork and many fees to meet and keep up with these requirements. We do not solicit donations in locations where we have not received written confirmation of compliance. To SEND DONATIONS or determine the status of compliance for any particular state visit [www.gutenberg.org/donate](http://www.gutenberg.org/donate)

While we cannot and do not solicit contributions from states where we have not met the solicitation requirements, we know of no prohibition against accepting unsolicited donations from donors in such states who approach us with offers to donate.

International donations are gratefully accepted, but we cannot make any statements concerning tax treatment of donations received from outside the United States. U.S. laws alone swamp our small staff.

Please check the Project Gutenberg Web pages for current donation methods and addresses. Donations are accepted in a number of other ways including checks, online payments and credit card donations. To donate, please visit: [www.gutenberg.org/donate](http://www.gutenberg.org/donate)

Section 5. General Information About Project Gutenberg-tm electronic works.

Professor Michael S. Hart was the originator of the Project Gutenberg-tm concept of a library of electronic works that could be freely shared with anyone. For forty years, he produced and distributed Project Gutenberg-tm eBooks with only a loose network of volunteer support.

Project Gutenberg-tm eBooks are often created from several printed editions, all of which are confirmed as Public Domain in the U.S. unless a copyright notice is included. Thus, we do not necessarily keep eBooks in compliance with any particular paper edition.

Most people start at our Web site which has the main PG search facility:

[www.gutenberg.org](http://www.gutenberg.org)

This Web site includes information about Project Gutenberg-tm, including how to make donations to the Project Gutenberg Literary Archive Foundation, how to help produce our new eBooks, and how to subscribe to our email newsletter to hear about new eBooks.